

Invited Paper

Chemicals from nature for weed management

Stephen O. Duke

Corresponding author. Natural Products Utilization Research Unit, Agricultural Research Service, United States Department of Agriculture, P.O. Box 8048, University, MS 38677; sduke@olemiss.edu

Franck E. Dayan

Agnes M. Rimando

Kevin K. Schrader

Natural Products Utilization Research Unit, Agricultural Research Service, United States Department of Agriculture, P.O. Box 8048, University, MS 38677

Giovanni Aliotta

Anna Oliva

Dipartimento di Scienze della Vita, Seconda Università degli Studi di Napoli, Via Vivaldi, 43-81100 Caserta, Italy

Joanne G. Romagni

Department of Biology, St. Thomas College, Houston, TX 77006

Natural products represent a vast repository of materials and compounds with evolved biological activity, including phytotoxicity. Some of these compounds can be used directly or as templates for herbicides. The molecular target sites of these compounds are often unique. Strategies for the discovery of these materials and compounds are outlined. Numerous examples of individual phytotoxins and crude preparations with weed management potential are provided. An example of research to find a natural product solution of a unique pest management problem (blue-green algae in aquaculture) is described. Finally, the problems associated with natural products for pest control are discussed.

Key words: Allelochemicals, aquaculture, blue-green algae, biological activity, natural products, phytotoxins.

Natural products have been the source of many pesticides, used either directly as crude preparations or as pure compounds. Rather than being used directly, they have been used more often as structural leads for the discovery and development of natural product-based pesticides. There are more examples of natural product use as fungicides, insecticides, and other pesticides than as herbicides (Pachlatko 1998). However, there are some success stories with herbicides, and natural products remain part of the herbicide discovery strategy for those companies that still have a herbicide discovery program.

The rationale for natural products in a herbicide discovery strategy has several disparate components. Nature is full of bioactive materials and compounds with unexploited properties. Many of the hundreds of thousands of secondary products generated by plants, microbes, and animals are the result of coevolution of the producing organism with pests. Thus, the compounds have biological activity. Sometimes, the function of the compound in nature is as a phytotoxin, as with phytotoxins produced by plant pathogens or allelochemicals produced by allelopathic plants. However, very often biocides with a specific function in nature can be used for quite different purposes, as with many natural product-based pharmaceuticals. In short, biological activity is more certain with secondary compounds from nature than with randomly synthesized compounds.

Improved instrumentation has considerably reduced the cost of isolation and identification of natural compounds from what it was a decade ago. This has caused renewed

interest in natural products in herbicide discovery programs. Another major reason for interest in natural phytotoxins is that they often have novel sites of action (Duke et al. 2000b, 2000d) (Table 1). Even if the phytotoxin is unsuitable for commercial use, identification of a new molecular target site can be very valuable in the design of synthetic herbicides. Natural compounds or preparations may require less regulatory scrutiny for registration than synthetic compounds, thus reducing the cost of commercializing the product. In some market niches, such as the home garden, the claim that a pesticide is “natural” will appeal to the consumer. In other markets, synthetic herbicides are either not allowed (organic gardening) or unlikely to have long-term approval (e.g., certain aquaculture situations). Lastly, despite the relatively low priority of natural products in herbicide discovery, there have been some major successes with natural products as herbicides or herbicide leads (described later). Successes lead to greater interest.

This review will discuss research strategies and the most important successes. Some interesting examples of natural products that have not made it to the market will also be provided, with the reasons that some of these products were unsuccessful. Several recent general reviews of natural products as potential herbicides are available (Dayan et al. 1999b; Duke et al. 1998, 2000b, 2000d, 2002; Hoagland 2001; Hoagland and Cutler 2000). Other reviews concentrate on microbial compounds as herbicides (Abbas and Duke 1997; Duke et al. 1996; Saxena and Pandey 2001) or the modes of action of natural phytotoxins (Duke et al.

TABLE 1. Molecular target sites of highly phytotoxic natural products.

Compound	Molecular target site ^a
AAL-toxin (a toxin from <i>Alternaria alternata</i>)	Ceramide synthase
Actinonin	Peptide deformylase
Brefeldin	Golgi apparatus function
Carbocyclic coformycin	Adenosine monophosphate deaminase
Cerulenin	3-ketoacyl-acyl-carrier protein synthase
Cochlioquinones	Mitochondrial reduced nicotinamide adenine dinucleotide reductase
Coronatine	Jasmonic acid antagonist
1,4-cineole	Asparagine synthetase
Fisherellin	<i>D-1 protein of photosystem II</i>
Fumonisin	Ceramide synthase
Fusicoccin	Plasma membrane adenosine triphosphatase (ATPase)
Gabaculin	Many transaminases
Gostatin	Aspartate amino transferase
Grandinol	<i>D-1 protein of photosystem II</i>
Hydantocidin	Adenylosuccinate synthase
Leptospermone	<i>4-Hydroxyphenylpyruvate dioxygenase</i>
Phaseolotoxin	Ornithine carbamoyltransferase
Phosphinothricin	Glutamine synthetase
Podophyllotoxin	<i>Tubulin</i>
Prehelminthosporol	Plasma membrane ATPase
Pyridazocidin	<i>Energy diversion from photosystem I</i>
Quassinoids	Nicotinamide adenine dinucleotide phosphate oxidase?
Rhizobitoxin	β-Cystathionase
Tagetitoxin	RNA polymerase
Sorgoleone	<i>D-1 protein of photosystem II</i>
Syringotoxin	Plasma membrane ATPase
Tentoxin	CF ₁ ATPase of the chloroplast
Tricolorin A	Plasma membrane ATPase
Thiolactomycin	Acetyl-CoA transacylase
Usnic acid	<i>4-Hydroxyphenylpyruvate dioxygenase</i>

^a Molecular target sites in italics are shared by synthetic herbicides.

1997, 2000c). Reviews of the general concept and strategies for utilizing natural products as pesticides are available (Rice et al. 1998). In this paper we shall update the material from these previous reviews and attempt to widen the scope of the coverage to include crude preparations and minor uses, such as natural products as algicides in aquaculture. However, this short review provides only a sampling of what is known about the potential use of natural products for weed management.

Discovery Strategies

Known Compounds

Hundreds of thousands of natural compounds have been isolated and their structures elucidated, but relatively few of these have been adequately tested for phytotoxicity. Rather than randomly bioassaying known compounds, structural and activity clues can be used to maximize chances of finding phytotoxic compounds.

Using Structural Clues

We know that certain chemical structures are more likely to affect a particular molecular target site than others. Thus, compounds with structures that are similar to the structures of known phytotoxins or inhibitors of particular plant enzymes or functions may have similar activities. For example, usnic acid (Figure 1), a secondary product from some li-

chens, is structurally similar to the triketone class of herbicides that inhibit 4-hydroxyphenylpyruvate dioxygenase (HPPD) (e.g., sulcotrione). This led Romagni et al. (2000b) to discover that (–)usnic acid is a better in vitro inhibitor of HPPD than sulcotrione. Natural products that are structural analogs of metabolic intermediates or enzyme cofactors might well be inhibitors of enzymes that use those intermediates or cofactors. For example, AAL-toxin (a toxin from *Alternaria alternata*) (Figure 1) is an analog of the sphingoid base substrates of ceramide synthase, and it is a potent inhibitor of this enzyme (Abbas et al. 1996, 2002). The sugar analog 2,5-anhydro-D-glucitol is phytotoxic (Figure 1) (Tanaka et al. 1996) because of its structural similarity to fructose (Dayan et al. 2002). This protoxin must be phosphorylated by the enzymes hexokinase and phosphofructokinase, yielding the bisphosphorylated analog of fructose-1,6-bisphosphate. Once bioactivated, this phosphorylated sugar analog inhibits fructose-1,6-bisphosphate aldolase.

Using Activity Clues

Many natural compounds have been tested for some type of biological activity other than herbicidal activity. If the mode of action of the activity is known, this information can be used to predict phytotoxicity and mode of action as a herbicide. For example, actinonin (Figure 1) and related compounds were known to be a potent new class of antibiotics with a unique mode of action, inhibition of peptide




FIGURE 1. Structures of some of the compounds produced by microbes mentioned in the text.

deformylase (Chen et al. 2000). This led Dirk et al. (2001) to find that their phytotoxicity is caused by the same mechanism.

Another example is that of the fumonisins (Figure 1). These compounds were first known to be toxic to animals by inhibiting ceramide synthase (Abbas et al. 1996). Fumonisins and structurally related compounds were found to be extremely phytotoxic by the same mechanism (Abbas et al. 1994).

Yet another example is that of the sesquiterpene lactone, artemisinin (Figure 2), from annual wormwood (*Artemisia annua* L.). This compound was isolated from this common plant because of ethnobotanical clues from Chinese folk medicine regarding its antimalarial properties. It is highly phytotoxic (Duke et al. 1987), but the mechanism of action of artemisinin and several active analogs is unknown (Dayan et al. 1999a). At least one company considered using artemisinin as the basis for development of a new herbicide. Antimalarial compounds might be expected to be herbicidal because *Plasmodium* spp. have a form of plastid with similarities to those of plants (Lang-Unnasch et al. 1998). Another natural product that is toxic to both plants (Zeidler et al. 1998) and *P. falciparum* (Jomaa et al. 1999) is fosmidomycin (Figure 1), an inhibitor of synthesis of all plastid terpenes. Conversely, the synthetic herbicide glyphosate has

been shown to be effective against *P. falciparum* (Roberts et al. 1999), and aryloxyphenoxypropionate herbicides are effective against the related parasite *Toxoplasma gondii* (Jelenska et al. 2001).

Discovery of New Compounds

Choosing a Biological Source

There are millions of organisms that have been inadequately studied for the biological properties of the secondary compounds that they produce. This is especially true for soil microbes. There are estimates that less than 1% of soil microbes have been cultured and identified (Felske et al. 1997; Pimm et al. 1995), leaving a potentially huge repository of unknown secondary compounds to be discovered. For the pesticide industry, soil microbes have been the organisms of choice for a source of natural products with potential for herbicides. Two strategies have been utilized: (1) isolation and culture of soil microbes from exotic locations, and (2) manipulation of culture conditions in order to culture previously uncultured organisms. Even when an organism can be cultured, there is no assurance that the culture conditions used will be adequate for the production of every secondary compound that the organism produces in nature.




FIGURE 2. Structures of some of the compounds produced by plants mentioned in the text.

Another strategy for phytotoxin discovery with microbes is the chemical ecology strategy. This approach is to examine only those organisms for which there is a reason to believe that a phytotoxin is produced. The most obvious case is that of plant pathogens (Sugawara 2000). For example, a dipeptide phytotoxin, maculosin (Figure 1), was found to be produced by the *Alternaria alternata* that infects spotted knapweed (*Centaurea maculosa* Lam.) (Bobylev et al. 1996). Many, if not most, plant pathogens produce phytotoxins to kill plant cells before they use them as a food source. The greatest difficulty of this approach has been problems in adequately culturing many plant pathogens.

A chemical ecology approach can also be taken in selecting plant species for extraction of phytotoxins. If a plant is known or suspected to be allelopathic, one should expect it to produce phytotoxic allelochemicals (Macias et al. 2001). There are many examples of this from the literature, although relatively few of these studies have adequately fractionated the plant for all phytotoxins. In many cases, bioassay-directed isolation was not carried out, and only water-soluble compounds were isolated.

Plants produce many highly phytotoxic compounds that have no obvious role in plant-plant interactions. For example, artemisinin is produced only in the glandular trichomes of annual wormwood (Duke et al. 1994), yet it is highly phytotoxic, as mentioned earlier. Hypericin (Figure 2), a highly phytotoxic compound, is produced only in certain cells of St. John's wort (*Hypericum perforatum* L.) (Duke et al. 2000a). Both of these compounds are toxic to the producing plant. In the case of artemisinin, the actual function of the compound in nature is unknown, whereas hypericin is an antifeedant. Plants sequester or secrete phytotoxic compounds to avoid autotoxicity (discussed in detail by Duke et al. 2000e). Thus, examination of plants and plant parts with specialized structures for sequestration of

secondary compounds is a good strategy for finding phytotoxins.

Bioassay-directed Isolation

Once a biological source is found, it must be fractionated into bioactive components and compounds (Duke et al. 2000b, 2000d, 2000e). Fractionation generally proceeds by initially partitioning extract from the identified biological source between polar and nonpolar solvents. Aliquots are tested for biological activity (e.g., phytotoxicity), and further fractionation is pursued on the fraction where activity is demonstrated. Column chromatography usually follows, and eluates (subfractions) are collected and tested. Chromatographic workup on the bioactive eluate (subfraction) is carried out until a pure compound is obtained. This whole process, which constitutes a bioassay-directed isolation, has resulted in many leads in the discovery of bioactive constituents from medicinal plants (e.g., Choudhary and Atta-ur-Rahman 1997; Pezzuto et al. 1998). Although still of limited practice in the discovery of pesticides, the isolation of an antifeedant compound and nematocidal alkaloids (Choudhary and Atta-ur-Rahman 1997), and phytotoxins from sorghum [*Sorghum bicolor* (L.) Moench] (Rimando et al. 1998), *Leucophyllum frutescens* (Rimando et al. 1999), and *Fusarium solani* (Tanaka et al. 1996) has been accomplished using this approach.

Some important things to consider when undertaking activity-guided natural product isolation are the choice of appropriate bioassay(s), extracting the plant part(s) where the bioactive constituent is most likely to be stored, and testing fractions and subfractions at equal concentrations. Following these guidelines, it was found that an allelopathic rice (*Oryza sativa* L.) cultivar 'Taichung Native 1' contained allelochemicals other than *p*-coumaric acid, a known phytotoxin

TABLE 2. Plant extracts that significantly inhibit germination and growth of the species tested. Data are arranged as follows: edible and medicinal plants are in alphabetical order, and weeds are listed according to their diffusion and economic importance.

Species, family, and biological form ^a	Affected plant(s)	References
Crops		
<i>Asparagus officinalis</i> L.; Liliaceae, Geophyte; asparagus	<i>Lycopersicon esculentum</i> Mill., <i>Asparagus officinalis</i> L., <i>Festuca</i> spp.	Weston 1996; Young 1986
<i>Brassica oleracea</i> L.; Cruciferae, Chamaephyte; cabbage	<i>Brassica oleracea</i> L., <i>Lactuca sativa</i> L., <i>Lycopersicon esculentum</i> Mill.	De Feo et al. 1997; Patrick and Koch 1958, 1963
<i>Canavalia ensiformis</i> (L.) DC.; Leguminosae, Therophyte; jack-bean	<i>Imperata brasiliensis</i> Trin.	Casini and Olivero 2001; Dinardo et al. 1998
<i>Crambe abyssinica</i> Hochst.; Cruciferae, Therophyte; crambe	<i>Triticum aestivum</i> L., <i>Abutilon theophrasti</i> Medicus	Vaughn and Berhow 1998
<i>Helianthus annuus</i> L.; Compositae, Therophyte; sunflower cultivars	<i>Amaranthus retroflexus</i> L., <i>Datura stramonium</i> L., <i>Hordeum vulgare</i> L., <i>Lactuca sativa</i> L., <i>Lepidium sativum</i> L., <i>Lycopersicon esculentum</i> Mill., <i>Triticum aestivum</i> L., <i>Trifolium</i> sp.	Barberi et al. 1998; Hall et al. 1982; Hilton 1979; Macias et al. 1996, 1999; Narwal et al. 1999
<i>Hordeum vulgare</i> L.; Gramineae, Therophyte; barley	<i>Sinapis alba</i> L.	Kati and Froud-Williams 1999
<i>Juglans nigra</i> L.; Juglandaceae, Phanerophyte; black walnut	<i>Lycopersicon esculentum</i> Mill.	Rice 1984; Willis 2000
<i>Olea europaea</i> L.; Oleaceae, Phanerophyte; olive oil mill wastewater	<i>Amaranthus retroflexus</i> L., <i>Chenopodium album</i> L., <i>Portulaca oleracea</i> L., <i>Raphanus sativus</i> L.	Aliotta et al. 2000
<i>Oryza sativa</i> L.; Gramineae, Therophyte; rice accessions	<i>Lactuca sativa</i> L., <i>Oryza sativa</i> L., <i>Echinochloa crus-galli</i> (L.) Beauv.	Chou and Lin 1976; Lin et al. 2000; Narwal 2000; Olofsdotter et al. 1999
<i>Sorghum bicolor</i> (L.) Moench; Gramineae, Therophyte; sorghum	<i>Festuca</i> spp.	Hall et al. 1982; Hilton 1979; Macias et al. 1996; Weston et al. 1999
<i>Secale cereale</i> L.; Gramineae, Therophyte; rye	<i>Chenopodium album</i> L.	Chou and Patrick 1976; Barnes et al. 1986
<i>Triticum aestivum</i> ; Gramineae, Therophyte; wheat accessions	Several weeds	Narwal 2000
<i>Ruta graveolens</i> L.; Rutaceae, Phanerophyte; rue	<i>Amaranthus retroflexus</i> L., <i>Chenopodium album</i> L., <i>Portulaca oleracea</i> L., <i>Raphanus sativus</i> L.	Aliotta and Cafiero 1999; Aliotta et al. 2000
<i>Zea mays</i> L.; Gramineae, Therophyte; maize	Several weeds	Bingamen and Christians 1995; Christians 1993; Gough and Carlstrom 1999
Weeds		
<i>Cyperus rotundus</i> L.; Cyperaceae, Geophyte; purple nutsedge	<i>Allium cepa</i> L., <i>Brassica nigra</i> (L.) Koch, <i>Brassica oleracea</i> L., <i>Cucumis sativus</i> L., <i>Daucus carota</i> L., <i>Fragaria</i> sp. pl., <i>Gossypium</i> sp., <i>Glycine max</i> (L.) Merr., <i>Hordeum vulgare</i> L., <i>Lycopersicon esculentum</i> Mill., <i>Oryza sativa</i> L., <i>Raphanus sativus</i> L., <i>Sorghum bicolor</i> (L.) Moench.	Friedman and Horowitz 1971; Lucena and Doll 1976; Singh 1968; Velu and Aruna 1996; Wibiwo 1996.
<i>Cynodon dactylon</i> (L.) Pers.; Gramineae, Geophyte; Bermuda grass	<i>Hordeum vulgare</i> L., <i>Glycine max</i> (L.) Merr.	Horowitz and Friedman 1971; Velu and Aruna 1996
<i>Echinochloa crus-galli</i> (L.) Beauv.; Gramineae, Therophyte; barnyard grass	<i>Triticum durum</i> L., <i>Glycine max</i> (L.) Merr., <i>Oryza sativa</i> L.	Dilday et al. 1998; Gressel and Holm 1964
<i>Eleusine indica</i> (L.) Gaertner; Gramineae, Therophyte; goosegrass	<i>Phaseolus vulgaris</i> L., <i>Sorghum bicolor</i> (L.) Moench, <i>Zea mays</i> L.	Altieri and Doll 1978
<i>Sorghum halepense</i> (L.) Pers.; Gramineae. Geophyte; johnson grass	<i>Coronilla varia</i> L., <i>Glycine max</i> (L.) Merr., <i>Gossypium</i> sp., <i>Hordeum vulgare</i> L.	Abdul-Wahab and Rice 1967; Bieber and Hoveland 1968; Horowitz and Friedman, 1971
<i>Imperata cylindrica</i> (L.) Beauv.; Gramineae, Geophyte; cogon grass	<i>Cucumis sativus</i> L., <i>Lycopersicon esculentum</i> Mill., <i>Oryza sativa</i> L., <i>Sorghum bicolor</i> (L.) Moench, <i>Zea mays</i> L.	Abdul-Wahab and Al-Naib 1972; Eussen 1978; Eussen and Soerjani 1978

TABLE 2. Continued.

Species, family, and biological form ^a	Affected plant(s)	References
<i>Portulaca oleracea</i> L.; Portulacaceae, Therophyte; purslane	<i>Medicago sativa</i> L., <i>Lycopersicon esculentum</i> Mill., <i>Triticum durum</i> L.	Le Tourmeau et al. 1956
<i>Chenopodium album</i> L.; Chenopodiaceae, Therophyte; lambs-quarter	<i>Beta vulgaris</i> L., <i>Brassica oleracea</i> L., <i>Triticum aestivum</i> L., <i>Zea mays</i> L.	Kossanel et al. 1977; Williams 1964
<i>Digitaria sanguinalis</i> (L.) Scop.; Gramineae, Therophyte; crabgrass	<i>Coronilla varia</i> L., <i>Harachis hypogea</i> L., <i>Gossypium</i> sp., Nitrificant bacteria	Gressel and Holm 1964; parenti and Rice 1969; Rice 1964
<i>Convolvulus arvensis</i> ; Convolvulaceae, Hemicytrophite; field bindweed	<i>Triticum aestivum</i> L. cv. Pavon	Alam et al. 1998
<i>Avena fatua</i> L.; Gramineae, Therophyte; wild oat	<i>Triticum aestivum</i> L., <i>Linum</i> sp., <i>Hordeum vulgare</i> L.	Schumacher et al. 1982; Tinnin and Muller 1971
<i>Amaranthus retroflexus</i> L.; Amaranthaceae, Therophyte; redroot pigweed	<i>Triticum durum</i> L., <i>Glycine max</i> (L.) Merr.	Bhowmik and Doll 1979; Qasem 1995

^a Chamaephyte: any perennial plant whose winter buds are within 25 cm of the soil surface. Geophyte: a perennial plant that is deeply embedded in the soil substrate. Hemicytrophite: a plant having buds at the soil surface and protected by scales, snow, or litter. Phanerophyte: a perennial tree or shrub with dormant buds borne on aerial shoots. Therophyte: an annual plant whose seed is the only overwintering structure.

found in the leaves and stems of rice (Chou et al. 1981). When tested at the same concentrations, the fraction containing *p*-coumaric acid, as well as a pure sample of *p*-coumaric acid, was not active against barnyardgrass [*Echinochloa crus-galli* (L.) Beauv.]. Phytotoxicity was detected in other fractions (Rimando et al. 2001).

Bioactivity-guided isolation is becoming an automated procedure where modern instrumentation in tandem is constructed to allow splitting of eluents, as peaks are detected, from extracts injected into a high-performance liquid chromatograph, a portion of which is collected into bioassay plates (e.g., 96-well plates), while another portion is diverted for structural analysis to a mass (Constant and Beecher 1995) or a nuclear magnetic resonance (NMR) spectrometer (Lindon et al. 1996). Fully automated analytical instrumentation, coupled with autosamplers for collecting fractions for bioassay, greatly reduces the time and effort put into the isolation of known compounds, while also maximizing the chance to identify new chemical entities with a specific biological activity (Hook et al. 1997). The power of a fully automated and integrated instrumentation has been demonstrated in the identification of 5-nitropyridone and other metabolites from corn (Bailey et al. 2000). Discovery of natural herbicides will be greatly facilitated using fully automated tandem instrumentations, such as liquid chromatography-mass spectrometry (LC/MS) and LC/MS/NMR.

Crude Extracts

There are a number of different ways to obtain collections of plants for novel drugs and allelochemical screening. These range from random screening to informed selection based on taxonomic, chemical, or ethnobotanical data, or any combination of these. As about three-quarters of the biologically active plant-derived compounds presently in use worldwide have been discovered through follow-up research to verify the authenticity of data from folk and ethnomedicinal uses (Farnsworth 1990), it is reasonable to conduct

ethnobotanically directed research in order to optimize the search for novel biologically active principles.

Since the emergence of agriculture and after a long period of adaptive coevolution between man and plants (Harlan 1992), an empirical body of knowledge has accumulated concerning the manipulation of ecosystems without the use of synthetic herbicides. Greek and Roman agronomists such as Theophrastus, Cato the Elder, Varro, Virgil, Columella, and Pliny the Elder wrote treatises on agriculture dealing with the methods of good crop husbandry used to minimize weed interference with crops, such as hand weeding, crop rotation, mechanical, tillage, burning, and mulching. Their insights came before any understanding of allelopathy or its mechanisms. Successively, farmers observed many allelopathy-related phenomena, for instance, that tomatoes (*Lycopersicon esculentum* L.) grow poorly under black walnut trees (*Juglans nigra* L.), and *sorghum* spp. plant residues and certain varieties of crop plants inhibit the growth of many weeds. We now know that many such relationships involve allelochemicals produced by plants (Waller 1987). Crude preparations of many plants significantly inhibit the germination and growth of species (Table 2). Theoretically, such preparations could be used as natural herbicides, although very little research has been conducted to exploit this approach. A more fruitful approach has been to use waste products from plants that are processed for food or oil.

For example, corn gluten meal, a by-product of the wet-milling process, is being used for preemergence weed management and fertilization (Bingamen and Christians 1995; Christians 1993; Gough and Carlstrom 1999). Corn gluten hydrolysate, produced by the action of a bacterial proteinase, is more active than the gluten meal as a herbicide (Liu and Christians 1994a). The hydrolysate is active on both grasses and broadleaf weeds (Liu and Christians 1997). The hydrolysate contains five phytotoxic dipeptides, with alanylalanine being the most active (Liu and Christians 1994b). Numerous physiological effects of alanylalanine have been reported

(Unruh et al. 1997a, 1997b), but the molecular target site is unknown. Later, a more active pentapeptide (leu-ser-pro-al-gln) was isolated from corn gluten hydrolysate (Liu and Christians 1996), but very little information is available on it.

Another seed meal waste product that could be used for weed management is that of crambe (*Crambe abyssinica* Hochst. ex R. E. Fries), a crop grown for oil and erucic acid (Vaughn and Berhow 1998). A major phytotoxin in this material is 1-cyano-2-hydroxy-3-butene (Figure 1). Seed meals from phytotoxic, glucosinolate-containing plants have also been shown to be potentially useful for weed management as a soil amendment (Vaughn et al. 1996).

Crude preparations of natural products must originate from crops or other organisms. There are many problematic issues with such products, such as supply, cost, and efficacy. The supply problem is likely to preclude widespread use in agronomic crops; however, certain niche markets could be ideal for this type of product.

Pure Compounds

Microbial Products

Products of Plant Pathogens

Plant pathogens produce a myriad of phytotoxins that are apparently useful in weakening the plant in the infection process. Most of these compounds are not host selective, even though the producing pathogen might infect only one or a limited number of host species. However, some of them are reported to be highly active against only one species or even certain genotypes of that species. Unless the host is a huge weed problem, there is little interest in herbicides that are extremely selective.

However, the literature on host-selective toxins is suspect from a herbicide standpoint. Plant pathologists have not always adequately evaluated these compounds for herbicidal activity. For example, AAL-toxin has been claimed to be a host-selective phytotoxin, affecting only certain tomato varieties (Abbas et al. 1995), yet it is one of the most generally phytotoxic natural products evaluated for herbicidal activity (Abbas et al. 1996). Other somewhat host-selective phytotoxins for weeds have been proposed, but not adequately tested. These include pyrenophorol (Figure 1) for wild oat (*Avena fatua* L.) control (Kastanias and Chrysai-Tokousbalides 2000) and maculosin (Figure 1) for spotted knapweed (Bobylov et al. 1996).

Space limitations do not permit mentioning many of the other phytotoxins from plant pathogens that have been described (Duke et al. 1996).

Other Microbial Compounds

Some of the more potent natural phytotoxins have come from nonpathogenic microbes. The bacterial phytotoxins of this type are reviewed by Barazani and Friedman (2001). We shall provide a few examples, with emphasis on those that have been most successful or spurred the most interest.

The actinomycete-produced compound, actinonin (also known as butanediamide) (Figure 1), is a potent inhibitor of peptide deformylase (Chen et al. 2000), an important enzyme of bacteria and chloroplasts (Dirk et al. 2001). Peptide deformylase is necessary for the posttranslational pro-

cessing of some chloroplast genome-encoded proteins. This is a site of action unique to plants and bacteria that has potential for the development of a new herbicide class. Although it is not extremely active on whole plants, it is highly active ($I_{50} < 100$ nM) on one out of two chloroplast peptide deformylases (Dirk et al. 2001).

Bialaphos (Figure 1) and phosphinothricin (Figure 1), both *Streptomyces* spp. products, are successful herbicides (Lydon and Duke 1999). Bialaphos is manufactured as a fermentation product and sold in a limited market in Japan. It is a proherbicide, requiring metabolic conversion to phosphinothricin in the target plant for herbicidal activity. Phosphinothricin is synthetically produced as glufosinate. It is the only herbicide that inhibits glutamine synthetase. It is nonselective, and many crops have been engineered to be resistant to it by inserting a transgene that encodes a detoxification gene from the producing *Streptomyces*. Phosphinothricin (glufosinate) is the biggest success story for a natural product-based herbicide. It is relatively inexpensive, toxicologically and environmentally safe, and efficacious on a wide range of target weeds. Thus, with herbicide-resistant crops, it has many of the same advantages of glyphosate in glyphosate-resistant crops.

Hydantocidin (Figure 1) is an *S. hygroscopicus* product with good herbicidal activity (reviewed by Duke et al. 1996, 2000b). It is a proherbicide that must be phosphorylated by the target plant to be an inhibitor of adenylosuccinate synthetase. Many analogs have been patented, but none of them have been marketed. Its nucleic acid synthesis site of action may be a cause for toxicological concern.

Pyridazocidin (Figure 1) is another *Streptomyces* sp. phytotoxin with an interesting mode of action (Gerwick et al. 1997). It is only a weak phytotoxin, but it is the only natural phytotoxin known to act by the same mechanism as paraquat, by accepting electrons from photosystem I and transferring them to molecular oxygen to generate a superoxide radical.

Cornexistin (Figure 1) and hydrocornexistin are products of a nonpathogenic fungus *Paecilomyces variotii*. Both of these compounds are very phytotoxic and have been patented as herbicides (Fields et al. 1996). Cornexistin has a unique molecular target site, aspartate amino transferase (Amagasa et al. 1994). The two compounds have different selectivity on crops and weeds.

New and interesting phytotoxins from microbes are being discovered at a high rate. For example, phomalactone (Figure 1) was recently found to cause rapid electrolyte leakage of plant plasma membranes, leading to plant death (Fukushima et al. 1998). Resormycin (Figure 1) (Igarashi et al. 1997) and the nigrosporins (Figure 1) (Tanaka et al. 1997) are interesting new phytotoxins with activity against both grasses and broadleaf weeds.

Lichens, a symbiotic relationship between algae and fungi, produce several different secondary compounds. The lichen is not a microbe, but the algae and fungi making it up could be considered microbes. Many lichen species are structurally unique, with ca. 50 to 60 of the 630 known lichen metabolites occurring in other fungi or higher plants. Most of the common lichen secondary metabolites are derived from the polyketide biosynthetic pathway (also called acetyl-polymalonyl pathway), with a few originating from the shikimate and mevalonate biosynthetic pathways. The

majority of lichen secondary metabolites originate from the fungal partner and are produced extracellularly when the organisms are in their obligate symbiotic association. Under specific conditions, where the symbionts have been artificially separated in the laboratory, other secondary compounds are produced; however, these are different from those produced in the symbiosis. Lichen secondary products may comprise up to 20% of the thallus dry weight, although 5 to 10% is more common. The unusual chemicals present in lichen and their biological activities can potentially be used to develop novel herbicide management tools.

The phytotoxic effect of certain lichen metabolites may deter the growth of other lichen populations as well as that of seedlings competing for a particular ecological niche. Several lichen species are found in the canopy and compete for space with ferns, fern allies, mosses, small seedlings, and other lichen species. Because lichens generally grow much slower than most of the competing species, they must have several defense strategies.

There is generally little information in the literature concerning the phytotoxic activity of specific lichen products. Most information concerns the antimicrobial or human health benefits of lichen secondary products. There are some studies, however, that document the specific activity of these compounds. The depsides barbatic acid and lecanoric acid (Figure 1), and the tridepside gyrophoric acid (Figure 1) act like PSII-inhibiting herbicides (e.g., atrazine, diuron) by interrupting photosynthetic electron transport in isolated chloroplasts (Endo et al. 1994; Rojas et al. 2000).

As mentioned previously, one interesting aspect of the allelopathic potential of lichens is related to the ability of (–) usnic acid to inhibit carotenoid biosynthesis through the enzyme HPPD (Romagni et al. 2000b). The *in vitro* activity of usnic acid is superior to that of other synthetic inhibitors of this herbicide target site. The ability of usnic acid to inhibit HPPD may contribute to an allelopathic effect. Synthetic inhibitors of this enzyme, e.g., sulcotrione and mesotrione, are currently being used as herbicides.

In another study, Lasceve and Gaugain (1990) documented the effects of usnic acid on sunflower (*Helianthus annuus* L.) and corn plants. There was a 40% decline in transpiration as well as dwarfism and root malformation. The actual mode of action was not determined; however, the concentration of usnic acid was 500 μM , and this might have caused secondary toxicity effects.

Abo-Khatwa et al. (1996) also studied three lichen acids, atranorin, usnic, and vulpinic acid, but found that they had characteristic effects similar to 2,4-dinitrophenol, an uncoupler of oxidative phosphorylation. The actual mode of action was not determined.

Several analogs of lichen-derived anthraquinones have strong herbicidal activity. Certain analogs of emodin, an anthraquinone found in nonlichen as well as lichenicolous fungi, are highly specific for grasses, causing malformation and bleaching in early seedlings. Other analogs of rhodocladonic acid cause root malformations in both dicot and monocot seedlings. Several of these completely inhibit germination.

Most of the lichen compounds are chemically simple, making them relatively easy to synthesize in the laboratory. Doing so would provide large amounts of material without harming the ecosystems. In addition, many of these compounds can be used as lead structures based on their partic-

ular mechanism of action and can then be optimized in the laboratory to fit specific applications.

Phytochemicals

Plants produce a very large number of interesting phytotoxins with potential use as herbicides. For example, leptospermone (Figure 2) is an allelochemical from which the triketone class of herbicides was developed (Mitchell et al. 2001). This is perhaps the most successful development of a commercial herbicide from a phytochemical.

Monoterpene cineoles are natural products commonly found in the essential oils from aromatic plants such as *Laurus nobilis* L., *Salvia* spp., *Eucalyptus* spp., and *Artemisia* spp. Many volatile monoterpenes are phytotoxic. Of those compounds, 1,8-cineole has been identified as one of the most potent allelochemicals released by *Artemisia* spp. 1,4-Cineole (Figure 2) is also common but is usually present in much lower concentrations. It has been recently reported that 1,4-cineole, a structural analog of the herbicide cinmethylin, was a potent inhibitor of asparagine synthetase (Romagni et al. 2000a).

Quassinoids are phytotoxic compounds produced by several plant species of the Simaroubaceae family (Heisey 1990; Lin et al. 1995). Chapparrinone (Figure 2) and its derivatives are active on plants at 1- to 5- μM concentrations. Chaparrinone-type quassinoids have broad weed spectrum activity when applied either as preemergence or postemergence herbicide and can provide 100% control of green foxtail [*Setaria viridis* (L.) Beauv.] and sicklepod [*Senna obtusifolia* (L.) Irwin & Barneby] at rates of 0.125 kg ha⁻¹. Quassinoids cause strong inhibition of the latter stages of mitosis but do not affect prophase, suggesting that these compounds do not prevent induction of the cell cycle. The primary site of action appears to be the inhibition of reduced nicotinamide adenine dinucleotide oxidase (Morré et al. 1998), and the presence of an oxymethylene bridge between C8 and C11 of the quassinoid backbone was required for the herbicidal activity (Dayan et al. 1999c).

There are many plant-derived compounds that inhibit mitosis of plants (Vaughan and Vaughn 1988). For example, podophyllotoxin (Figure 2) is an aryltetralin lignan extracted from the leaves of mayapple (*Podophyllum peltatum* L.) and has been studied extensively because of its ability to inhibit microtubule assembly in mammalian cells and its strong antiviral activity (Imbert 1998). Numerous semisynthetic derivatives of podophyllotoxin, such as etoposide and teniposide, having inhibitory activity against DNA-topoisomerase II, have been developed as effective antineoplastic drugs (Canel et al. 2000). When tested for phytotoxicity, this compound was more active against the monocotyledonous Italian ryegrass (*Lolium multiflorum* Lam.) than against the dicotyledonous lettuce (*Lactuca sativa* L.) species tested (Oliva et al. 2002). Moreover, the inhibition of root growth observed in the bioassays suggested that podophyllotoxin affected either cell division or cell elongation. Its inhibitory effect on the different mitotic phases, and especially on prophase, indicates that this compound affects mitosis. In addition, abnormal mitotic stages observed in root cells exposed to 100 μM podophyllotoxin suggested that this compound acted as a natural mitotic disrupter (Vaughan and Vaughn 1990). Effects were noted only on the cell division stages past prometaphase. At this stage, the normal spindle

microtubular organizing center seemed to be affected by the tested compound, so that the microtubules were oriented toward multiple poles where the chromatids appeared in "star anaphase" figures. This display is similar to that found when the herbicides terbutol, sindone B, pronamide, di-thiopyr, and some carbamate herbicides were tested at low concentrations (0.1 to 30 μM) (Hoffman and Vaughn 1994; Lehnert and Vaughn 1992; Lehnert et al. 1990). Although the precise molecular mechanism of action of this compound remains to be identified in plants, a primary effect is the alteration of the formation of the spindle microtubular organization centers, resulting in the formation of multiple spindle poles, leading to an asymmetrical convergence of the chromatids.

Sorgoleone (Figure 2) is an allelochemical exuded in oily droplets from the roots of sorghum (Nettly and Butler 1986). The concentration of sorgoleone in soils growing sorghum can reach 10^{-4} to 10^{-5} M (Nettly et al. 1988) and lead to suppression of weed growth (Forney and Foy 1985). This potent phytotoxin represses the growth of large crabgrass seedlings [*Digitaria sanguinalis* (L.) Scop.] with a GR_{50} of 10 μM for shoot and root growth (Nimbal et al. 1996). Inhibition of shoot and root growth of velvetleaf (*Abutilon theophrasti* Medikus) and barnyardgrass have also been observed at concentrations from 10 to 200 μM . Sorgoleone inhibits photosynthetic electron transport in thylakoids by competing for the plastoquinone binding site on photosystem II. This compound has the same level of inhibitory activity of photosynthetic electron transport as diuron, 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU), with an I_{50} of 100 and 120 nM, respectively (Gonzalez et al. 1997; Nimbal et al. 1996).

Multicolored morningglory (*Ipomoea tricolor* Cav.) is used in traditional agriculture in Mexico for weed management. It contains resin glycosides of which tricolorin A (Figure 2) is the principal constituent (Pereda-Miranda et al. 1993). This compound is highly phytotoxic and a potent inhibitor of plasma membrane adenosine triphosphatase (Calera et al. 1995).

Other Phytochemicals

Grandinol (Figure 2) is a phloroglucinol-related compound that inhibits germination, photosynthesis, transpiration, and stomatal opening (Yoneyama et al. 1996).

Natural Compounds from Other Sources

Secondary products with many kinds of biological activity are produced by all types of marine organisms and animals. For example, certain ants apparently produce herbicidal compounds to kill the vegetation surrounding their habitat (Renner and Ricklefs 1998). Relatively little effort has been made to examine such compounds for their potential as herbicides.

Natural Compounds for Use as Algicides—A Niche Market

Aquaculture is one of the fastest growing areas of agriculture in the world, and in the United States catfish is the largest and fastest growing segment of aquaculture. Off-flavors, which render cultured catfish unpalatable and thereby

unmarketable, cause large economic losses every year. Certain species of cyanobacteria (blue-green algae) that grow in catfish aquaculture ponds produce compounds that cause objectionable flavors. The musty flavor of 2-methylisoborneol (MIB) is most commonly found in the flesh of cultured catfish raised in Mississippi, which provides over half of the world's supply of farm-raised channel catfish (*Ictalurus punctatus*). The cyanobacterium *Oscillatoria perornata* (Skuja) has been identified as the leading cause of the musty off-flavor in catfish produced in west Mississippi (van der Ploeg et al. 1992).

The only compounds approved by the United States Environmental Protection Agency (USEPA) for application as an algicide in catfish production ponds in Mississippi are diuron and copper-based products, such as chelated-copper products and copper sulfate. Currently, the USEPA must grant catfish farmers an emergency exemption for every year that it is to be used, and therefore there is no guarantee of renewal each year. The algicides currently used are not selective and kill other groups of phytoplankton, such as green algae (Chlorophyta) (Schrader et al. 1998a). Green algae are the preferred type of phytoplankton, compared with cyanobacteria, in catfish production (Paerl and Tucker 1995). Additional undesirable attributes of the use of synthetic compounds as algicides in aquaculture ponds include environmental safety issues and the public's negative perception of the use of synthetic chemicals in food-production ponds.

The discovery of environmentally safe natural compounds that are selectively toxic toward odor-producing species of cyanobacteria would greatly benefit commercial catfish producers. There are few studies concerning the use of natural compounds as selective algicides in catfish ponds. The earliest reported studies involve the discovery of several oxygenated fatty acids (allelochemicals produced by the aquatic plant *Eleocharis microcarpa*) that inhibited the growth of cyanobacteria found in catfish ponds (Van Aller and Pessoney 1982; Van Aller et al. 1985). Potassium ricinoleate was determined to be structurally similar to the active fatty acids, and the commercial algicide Solricin 135[®] (Caschem, Inc., Bayonne, NJ) contains potassium ricinoleate as the active ingredient. However, Tucker and Lloyd (1987) found that potassium ricinoleate was ineffective in controlling undesirable species of cyanobacteria in efficacy studies conducted in catfish ponds. Another more recent study has evaluated the use of decomposing barley (*Hordeum vulgare* L.) straw to control cyanobacteria in catfish ponds (Wills et al. 1999). Welch et al. (1990) were among the first to report the reduction of filamentous algae because of the presence of decomposing barley straw placed in a canal in the U.K. One possible mechanism for algal inhibition may be the result of the release of antialgal compounds from the decomposing barley straw (Pillinger et al. 1994). Studies by Wills et al. (1999) have found that decomposing barley straw does not prevent musty off-flavor in cultured catfish.

Recent discoveries by Schrader and Harries (2001) and Schrader et al. (1998b) have identified several promising natural compounds produced by certain plants that are selectively toxic toward *O. perornata*. One of these compounds, ferulic acid (Figure 2), was pond-tested, and numbers of *O. perornata* were not significantly reduced by ferulic acid application (Schrader et al. 2000b). These unfavorable results were attributed to the rapid dissipation of ferulic acid

from the water column. This study provides an example of a potential concern for using natural compounds as algicides in highly eutrophic catfish aquaculture ponds. In such an ecosystem certain toxic natural compounds may undergo rapid dissipation, degradation, or transformation, thereby limiting sufficient contact with or uptake by the target microbe to provide an effective means for elimination of the noxious organism.

Anthraquinone (Figure 2) is another promising natural compound identified as a selective algicide during laboratory screening (Schrader et al. 1998b). Anthraquinone appears to inhibit electron transfer in photosystem II in *O. perornata* (Schrader et al. 2000a). Subsequent efficacy testing of anthraquinone found that it was ineffective in reducing the numbers of *O. perornata* or the levels of MIB in pond water when applied in a solution of ethanol or as an oil emulsion (Schrader, unpublished data). Anthraquinone ineffectiveness as a selective algicide in pond tests was attributed to its lack of solubility in water. This problem highlights another consideration in using a natural compound as a selective algicide in that the compound must be in a form or applied in a way that maintains it in the water in an active (toxic) state so that it can effectively enter the target organism.

There are also considerations in selecting natural compound leads for subsequent development as commercial algicides. For example, the cost to the catfish farmer must be considered. A compound must be effective in controlling the undesirable cyanobacterial species at a low concentration ($\mu\text{g L}^{-1}$ range), and the cost of synthesis at the industrial-scale level will affect the cost to the farmer. Also, there are several environmental safety concerns that must be addressed, including persistence of the natural compound in the catfish pond water and sediment, potential breakdown products of the compound, its toxicity toward nontarget organisms (e.g., catfish), and its potential accumulation in and metabolism by catfish.

Limitations and Future Research

We have described the benefits and successes of using natural products as herbicides or for herbicide discovery. So, why is there not more interest in this approach? There are a number of problems associated with natural products that have deterred a stronger interest.

First, many natural products are too expensive to be seriously considered for use as agrochemicals because of their structural complexity. For example, the cyclic tetrapeptide rentoxin (Figure 1) is an excellent herbicide but is very expensive. Considerable structure-activity research to find a simpler, less expensive molecule with similar activity (e.g., Bland et al. 1993; Edwards et al. 1988) was unsuccessful. Conversion of the molecule to another complex structure yields isotentoxin, a compound with a different mode of action (Liebermann et al. 1996). Another example is that of carbocyclic coformycin (Pillmoor 1998) (Figure 1). This compound with a unique mode of action (Table 1) could not be synthesized economically. Even close analogs had little or no herbicidal activity. In cases where the function of the compound in nature is that of a phytotoxin, the structure may have been optimized by nature. In such cases, if the structure is complex, attempts to generate a low-cost molecule may be unsuccessful.

The popular media bombard the public with the medieval herbalist concept that everything in nature is healthy, but many of the most toxic compounds known to humans are natural (e.g., aflatoxin, fumonisins, ricin). Some highly phytotoxic natural products are also very toxic to mammals. For example, AAL-toxin is quite toxic to mammalian cells (Abbas et al. 1996). This aspect of some natural phytotoxins has terminated interest in them for the development for weed management. Nevertheless, from an environmental toxicology standpoint, the relatively short half-life of most natural compounds in the field is desirable. Regulation of natural product-based pesticides varies from one country to another (Neale 2000). According to current regulations, the USEPA may require the same scrutiny for a natural product as for a synthetic pesticide.

The intellectual property rights associated with rare organisms from exotic places have become quite complicated. Developing countries, especially, have felt that their biological resources have been exploited by institutions from the developed world for the discovery of pharmaceuticals. This has resulted in these countries passing laws to retain some level of ownership of compounds discovered from organisms taken from within their borders (ten Kate and Laird 1999). In some cases, this has discouraged discovery efforts that use the biological diversity of certain places.

Until recently, the cost of compound isolation and structure elucidation was very high. This problem, coupled with the rediscovery of known compounds (e.g., Ayer et al. 1989; Heisey et al. 1988), was discouraging. Modern tandem instrumentation, such as LC/MS/MS and LC/NMR, and automation of determination of compounds within a complex extract (Hook et al. 1997) has greatly reduced the time and effort needed to identify isolated compounds. Precise profiling of known compounds in standard bioassays and use of this information to eliminate costly purification and structure elucidation of known compounds has been a strategy to reduce cost. However, if standards are set too high, this approach will result in the loss of some new compounds with similar activity to known compounds. Tandem analytical instrumentation should eliminate this choice, allowing rapid and inexpensive determination of whether the suspected known compound is in the fraction. If not, bioassay-directed isolation can proceed.

The relatively short environmental half-life of natural products is good from an environmental toxicology standpoint, but a herbicide must persist sufficiently long to have the desired effect. We have found, in some cases, that some natural herbicides do not persist long enough to be effective (e.g., Schrader et al. 2000b).

Synthetic chemistry is the dominant approach to herbicide discovery. The recent high level of investment in combinatorial chemical synthesis and high throughput screening has reinforced this approach. Pillmoor (1998) stated in his discussion of natural product approaches that "unless the commitment is there to invest in this work, then the value of undertaking natural product research in the first place is somewhat questionable." We believe that modern instrumentation and improved methods should increase interest in natural product-based herbicide discovery research.

Literature Cited

Abbas, H. K. and S. O. Duke. 1997. Plant pathogens and their phytotoxins as herbicides. Pages 1–20 in R. K. Upadhyay and K. G. Mukerji, eds.

- Toxins in Plant Disease Development and Evolving Biotechnology. New Delhi: Oxford & IBH Publishing.
- Abbas, H. K., S. O. Duke, W. T. Shier, and M. V. Duke. 2002. Inhibition of ceramide synthesis in plants by phytotoxins. In R. K. Upadhyay, ed. *Advances in Microbial Toxin Research and Its Biotechnological Exploitation*. London: Kluwer Academic Publishers/Plenum Press. In press.
- Abbas, H. K., S. O. Duke, W. T. Shier, R. T. Riley, and G. A. Kraus. 1996. The chemistry and biological activities of the natural products AAL-toxin and the fumonisins. Pages 293–308 in B. R. Singh and A. T. Tu, eds. *Natural Toxins 2. Structure, Mechanism of Action, and Detection*. *Advances in Experimental Medicine and Biology*. Volume 391. New York: Plenum Press.
- Abbas, H. K., T. Tanaka, S. O. Duke, et al. 1994. Fumonisin and AAL-toxin-induced disruption of sphingolipid metabolism with accumulation of free sphingoid bases: involvement in plant disease. *Plant Physiol.* 106:1085–1093.
- Abbas, H. K., T. Tanaka, S. O. Duke, and C. D. Boyette. 1995. Susceptibility of various crop and weed species to AAL-toxin, a natural herbicide. *Weed Technol.* 9:125–130.
- Abdul-Wahab, A. S. and F. A. G. Al-Naib. 1972. Inhibitional effects of *Imperata cylindrica* (L.) P. B. Bull. Iraq Nat. Hist. Mus. 5:17–24.
- Abdul-Wahab, A. S. and E. L. Rice. 1967. Plant inhibition by Johnson grass and its possible significance in old field succession. Bull. Torrey Bot. Club 94:486–497.
- Abo-Khatwa, A. N., A. A. Al-Robai, and D. A. Al-Jawhari. 1996. Lichen acids as uncouplers of oxidative phosphorylation of mouse-liver mitochondria. *Nat. Toxins* 4:96–102.
- Alam, S. M., A. R. Azmi, S. A. Alam, S. S. M. Naqvi, and R. Ansari. 1998. Effect of aqueous leaf extract of field bindweed (*Convolvulus arvensis* L.) and salinity on growth of wheat. *Rachis* 17:49–51.
- Aliotta, G. and G. Cafiero. 1999. Biological properties of *Ruta graveolens* L. and its potential use in sustainable agricultural systems. Pages 551–563 in Inderjit, K.M.M. Dakshiny, and C. L. Foy, eds., *Principles and Practice in Plant Ecology*. Boca Raton, FL: CRC Press.
- Aliotta, G., G. Cafiero, V. De Feo, B. Di Blasio, R. Iacovino, and A. Oliva. 2000. Allelochemicals from rue (*Ruta graveolens* L.) and olive (*Olea europaea* L.) oil mill waste waters as potential natural pesticides. *Curr. Top. Phytochem.* 3:167–177.
- Altieri, M. A. and J. D. Doll. 1978. The potential of allelopathy as a tool for weed management in crops. *PANS* 24:495–502.
- Amagasa, T., R. N. Paul, J. J. Heitholt, and S. O. Duke. 1994. Physiological effects of cornexistin on *Lemna paucicostata*. *Pestic. Biochem. Physiol.* 49:37–52.
- Ayer, S. W., B. G. Isaac, D. M. Krupa, K. E. Crosby, L. J. Letendre, and R. J. Stonard. 1989. Herbicidal compounds from microorganisms. *Pestic. Sci.* 27:221–223.
- Bailey, N. J. C., P. D. Stanley, S. T. Hadfield, J. C. Lindon, and J. K. Nicholson. 2000. Mass spectrometrically detected directly coupled high performance liquid chromatography/nuclear magnetic resonance spectroscopy/mass spectrometry for the identification of xenobiotics metabolites in maize plants. *Rapid Commun. Mass Spectrom.* 14: 679–684.
- Barazani, O. and J. Friedman. 2001. Allelopathic bacteria and their impact on higher plants. *Crit. Rev. Microbiol.* 27:41–55.
- Barberi, P., A. Cozzani, M. Macchia, and E. Bonari. 1998. Size and composition of the weed seedbank under different management systems for continuous maize cropping. *Weed Res.* 38:319–334.
- Barnes, J. P., A. R. Putnam, and B. A. Burke. 1986. Allelopathic activity of rye (*Secale cereale* L.). Pages 271–286 in A. Putnam and C. S. Tang, eds. *The Science of Allelopathy*. New York: Wiley Interscience.
- Bhowmik, P. C. and J. D. Doll. 1979. Evaluation of allelopathic effects of selected weed species on corn and soybeans. *Proc. N. Cent. Weed Control Conf.* 34:43–45.
- Bieber, G. L. and C. S. Hoveland. 1968. Phytotoxicity of plant materials on seed germination of crownvetch, *Coronilla varia* L. *Agron. J.* 60: 185–188.
- Bingamen, B. R. and N. E. Christians. 1995. Greenhouse screening of corn gluten meal as a natural control product for broadleaf and grassy weeds. *HortSci.* 30:1256–1259.
- Bland, J. M., J. V. Edwards, S. R. Eaton, and A. R. Lax. 1993. Potential of natural peptide compounds as leads for novel pesticides. *Pestic. Sci.* 39:331–340.
- Bobylev, M. M., L. I. Bobyleva, and G. A. Strobel. 1996. Synthesis and bioactivity of analogs of maculosin, a host-specific phytotoxin produced by *Alternaria alternata* on spotted knapweed (*Centaurea maculosa*). *J. Agric. Food Chem.* 44:3940–3964.
- Calera, M. R., A. L. Anaya, and M. Gavilanes-Ruiz. 1995. Effect of phytotoxic resin glycoside on activity of H⁺-ATPase from plasma membrane. *J. Chem. Ecol.* 21:289–297.
- Canel, C., R. M. Moraes, F. E. Dayan, and D. Ferreira. 2000. Podophyllotoxin. *Phytochemistry* 54:115–120.
- Casini, P. and L. Olivero. 2001. Allelopathic effects of legume cover crops on cogon grass (*Imperata brasiliensis* Trin.). *Allelopathy J.* 88:189–200.
- Chen, D. Z., D. V. Patel, C. J. Hackbarth, et al. 2000. Actinonin, a naturally occurring antibacterial agent, is a potent deformylase inhibitor. *Biochemistry* 39:1256–1262.
- Chou, C. H., Y. C. Chiang, and H. H. Cheng. 1981. Autointoxication mechanism of *Oryza sativa*. III. Effect of temperature on phytotoxin production during rice straw decomposition in soil. *J. Chem. Ecol.* 7: 56–67.
- Chou, C. H. and H. J. Lin. 1976. Autointoxication mechanisms of *Oryza sativa*. I. Phytotoxic effects of decomposing rice residues in soil. *J. Chem. Ecol.* 2:353–367.
- Chou, C. H. and Z. A. Patrick. 1976. Identification and phytotoxic activity of compounds produced during decomposition of corn and rye residues in soil. *J. Chem. Ecol.* 2:369–387.
- Choudhary, M. I. and Atta-ur-Rahman. 1997. Bioactivity-guided phytochemicals from medicinal plants. *R. Soc. Chem. Spec. Publ.* 200(Phytochem. Diversity):41–52.
- Christians, N. E. 1993. The use of corn gluten meal as a natural pre-emergence weed control in turf. Pages 284–290 in R. N. Carrow, N. E. Christians, and R. C. Shearman, eds. *International Turfgrass Society Research Journal*. Overland Park, KS: Intertec Publishing.
- Constant, H. and C.W.W. Beecher. 1995. A method for the dereplication of natural product extracts using electrospray HPLC/MS. *Nat. Prod. Lett.* 6:193–196.
- Dayan, F. E., A. Hernandez, S. N. Allen, R. M. Moraes, J. A. Vroman, M. A. Avery, and S. O. Duke. 1999a. Comparative phytotoxicity of artemisinin and several sesquiterpene analogues. *Phytochemistry* 50: 607–614.
- Dayan, F. E., A. M. Rimando, M. R. Tellez, B. E. Scheffler, T. Roy, and H. K. Abbas. 2002. The mechanism of action of the fungal phytotoxin 2,5-anhydro-D-glucitol requires bioactivation by glycolytic enzymes. *Weed Sci. Soc. Am. Abstr.* 42:49.
- Dayan, F. E., J. G. Romagni, M. R. Tellez, A. M. Rimando, and S. O. Duke. 1999b. Managing weeds with natural products. *Pestic. Outlook* 10:185–188.
- Dayan, F. E., S. B. Watson, J.C.G. Galindo, A. Hernandez, J. Dou, J. D. McChesney, and S. O. Duke. 1999c. Phytotoxicity of quassinoids: physiological responses and structural requirements. *Pestic. Biochem. Physiol.* 65:15–24.
- De Feo, V., F. De Simone, F. Giannattasio, V. Magnifico, A. Marcantonio, and A. D. Palumbo. 1997. Allelopathic effects of broccoli (*Brassica oleracea* L. var. italica Plenck) extract on vegetable crops. *Allelopathy J.* 4:277–281.
- Dilday, R. H., K. A. Moldenhauer, W. G. Yan, and D. R. Gealy. 1998. Allelopathic activities of barnyard grass in rice and yield reduction due to barnyard grass infestation. *Ark. Agric. Exp. Stn.* 460:27–31.
- Dinardo, W., M. T. Pellegrini, and P. L. C. A. Alves. 1998. Inhibitory effects of Jackbean (*Canavalia ensiformis* L.) leaf residue on germination and vigour of crops and weeds. *Allelopathy J.* 5:35–42.
- Dirk, L. M. A., M. A. Williams, and R. L. Houtz. 2001. Eukaryotic peptide deformylases, nuclear-encoded and chloroplast-targeted enzymes in *Arabidopsis*. *Plant Physiol.* 127:97–107.
- Duke, S. O., H. K. Abbas, T. Amagasa, and T. Tanaka. 1996. Phytotoxins of microbial origin with potential for use as herbicides. Pages 82–113 in L. G. Copping, ed. *Crop Protection Agents from Nature: Natural Products and Analogues*. *Critical Reviews on Applied Chemistry*. Volume 35. Cambridge, U.K.: Society for Chemical Industries.
- Duke, S. O., S. R. Baerson, F. E. Dayan, I. A. Kagan, A. Michel, and B. E. Scheffler. 2001. Biocontrol of weeds without the biocontrol agent. Pages 96–105 in J. Gressel and G. M. Vurro, eds. *NATO Advanced Research Workshop: Enhancing Biocontrol Agents and Handling Risks*. Amsterdam: IOS Press.
- Duke, S. O., C. Canel, A. M. Rimando, M. R. Tellez, M. V. Duke, and R. N. Paul. 2000a. Current and potential exploitation of plant glandular trichome productivity. *Adv. Bot. Res.* 31:121–151.
- Duke, S. O., F. E. Dayan, A. Hernandez, M. V. Duke, and H. K. Abbas. 1997. Natural products as leads for new herbicide modes of action. *Brighton Crop Prot. Conf., Weeds—1997.* 2:579–586.

- Duke, S. O., F. E. Dayan, and A. M. Rimando. 1998. Natural products as tools for weed management. *Proc. Jpn. Weed Sci. Soc. (Suppl.)*:1–11.
- Duke, S. O., F. E. Dayan, and A. M. Rimando. 2000b. Natural products and herbicide discovery. Pages 105–133 in A. H. Cobb and R. C. Kirkwood, eds. *Herbicides and their Mechanisms of Action*. Sheffield, U.K.: Academic Press.
- Duke, S. O., F. E. Dayan, and J. G. Romagni. 2000c. Natural products as sources for new mechanisms of herbicidal action. *Crop Prot.* 19:583–589.
- Duke, S. O., F. E. Dayan, J. G. Romagni, and A. M. Rimando. 2000d. Natural products as sources of herbicides: current status and future trends. *Weed Res.* 40:99–111.
- Duke, M. V., R. N. Paul, H. K. Elsohly, G. Sturtz, and S. O. Duke. 1994. Localization of artemisinin and artemisitene in foliar tissues of glanded and glandless biotypes of *Artemisia annua*. *Int. J. Plant Sci.* 155:365–373.
- Duke, S. O., A. M. Rimando, F. E. Dayan, et al. 2000e. Pages 1–20 in W. R. Bidlack, S. T. Omaye, M. S. Meskin, and D.K.W. Topham, eds. *Strategies for the Discovery of Bioactive Phytochemicals*. Phytochemicals as Bioactive Agents. Lancaster, PA: Technomic Publishing.
- Duke, S. O., K. C. Vaughn, E. M. Croom, Jr., and H. N. Elsohly. 1987. Artemisinin, a constituent of annual wormwood (*Artemisia annua*), is a selective phytotoxin. *Weed Sci.* 35:499–505.
- Edwards, J. V., O. D. Dailey, J. M. Bland, and H. G. Cutler. 1988. Approaches to structure–function relationships for naturally occurring cyclic peptides. *Am. Chem. Soc. Symp. Ser.* 380:35–56.
- Endo, Y., H. Hayashi, T. Sato, M. Maruno, T. Ohta, and S. Nozoe. 1994. Confluent acid and 2'-O-methylperlatolic acid, monoamine oxidase B inhibitors in a Brazilian plant, *Himatanthus sucuuba*. *Chem. Pharm. Bull.* 42:1198–1201.
- Eussen, J. H. H. 1978. Isolation of growth inhibiting substances from alang-alang (*Imperata cylindrica* (L.) Beauv.). In J.H.H. Eussen, ed. *Studies on the Tropical Weed Imperata cylindrica* (L.) var. major. Paper No. 7. Utrecht, The Netherlands: Drukkerij Elinkwijk BV.
- Eussen, J. H. H. and M. Soerjani. 1978. Allelopathic activity of alang-alang (*Imperata cylindrica* (L.) Beauv.), isolation of growth regulating substances from leaves. In J.H.H. Eussen, ed. *Studies on the Tropical Weed Imperata cylindrica* (L.) Beauv. var. major. Paper No. 6. Utrecht, The Netherlands: Drukkerij Elinkwijk BV.
- Farnsworth, N. R. 1990. The role of ethnopharmacology and the search for new drugs. Pages 2–21 in J. Chadwick and J. Marsh, eds. *Bioactive Compounds from Plants*. Chichester, U.K.: J. Wiley.
- Felske, A., H. Rheims, A. Wolternick, E. Stackebrandt, and A. D. Akkermans. 1997. Ribosome analysis reveals prominent activity of an uncultured member of the class Actinobacteria in grassland soils. *Microbiology* 143:2983–2989.
- Fields, S. C., L. Mireles-Lo, and B. C. Gerwick. 1996. Hydroxycornexistin: a new phytotoxin from *Paecilomyces variotii*. *J. Nat. Prod.* 59:698–700.
- Forney, D. R. and C. L. Foy. 1985. Phytotoxicity of products from rhizospheres of a sorghum–sudangrass hybrid (*S. bicolor* × *S. sudanese*). *Weed Sci.* 33:597–604.
- Friedman, T. and M. Horowitz. 1971. Biologically active substances in subterranean parts of purple nutsedge. *Weed Sci.* 19:398–401.
- Fukushima, T., M. Tanaka, M. Gohbara, and T. Fujimori. 1998. Phytotoxicity of three lactones from *Nigrospora sacchari*. *Phytochemistry* 48: 625–630.
- Gerwick, B. C., S. S. Fields, P. R. Graupner, J. A. Gray, E. L. Chapin, J. A. Cleveland, and D. R. Heim. 1997. Pyridazocidin, a new microbial phytotoxin with activity in the Mehler reaction. *Weed Sci.* 45:654–657.
- Gonzalez, V., C. I. Nimbal, L. A. Weston, and G. M. Cheniae. 1997. Inhibition of a photosystem II electron transfer reaction by sorgoleone, a natural product. *J. Agric. Food Chem.* 45:1415–1421.
- Gough, R. E. and R. Carlstrom. 1999. Wheat gluten meal inhibits germination and growth of broadleaf and grassy weeds. *HortSci.* 34:269–270.
- Gressel, J. B. and L. G. Holm. 1964. Chemical inhibition of crop germination by weed seeds and the nature of inhibition by *Abutilon theophrasti*. *Weed Res.* 4:44–53.
- Hall, A. B., U. Blum, and R. C. Fites. 1982. Stress modification of allelopathy of *Helianthus annuus* L. debris on seed germination. *Am. J. Bot.* 69:776–783.
- Harlan, J. 1992. *Crops and Man*. Madison, WI: American Society of Agronomy and American Crop Science Society. p. 41.
- Heisey, R. 1990. Allelopathic and herbicidal effects of extracts from tree of heaven (*Ailanthus altissima*) Am. J. Bot. 77:662–670.
- Heisey, R. M., S. K. Mishra, and A. R. Putnam. 1988. Production of herbicidal and insecticidal metabolites by soil microorganisms. *Am. Chem. Soc. Symp. Ser.* 380:68–78.
- Hilton, J. L. 1979. Research on the physiology and biology of weeds. *Weeds Today* 10(4):5–6.
- Hoagland, R. E. 2001. Bioherbicides: phytotoxic natural products. *Amer. Chem. Soc. Symp. Ser.* 774:72–90.
- Hoagland, R. E. and S. J. Cutler. 2000. Plant and microbial compounds as herbicides. Pages 73–99 in S. S. Narwal, R. E. Hoagland, R. H. Dilday, and M. J. Reigosa, eds. *Allelopathy in Ecological Agriculture and Forestry*. Amsterdam: Kluwer Academic Publishers.
- Hoffman, J. C. and K. C. Vaughn. 1994. Mitotic disrupter herbicides act by a single mechanism but vary in efficacy. *Protoplasma* 179:16–25.
- Hook, D. J., E. J. Pack, J. J. Yacobucci, and J. Guss. 1997. Approaches to automating the dereplication of bioactive natural products—the key step in high throughput screening of bioactive materials from natural sources. *J. Biomol. Screening* 2:145–152.
- Horowitz, M. and T. Friedman. 1971. Biological activity of subterranean residues of *Cynodon dactylon* L., *Sorghum halepense* L., and *Cyperus rotundus* L. *Weed Res.* 11:88–93.
- Igarashi, M., N. Kinoshita, T. Ikeda, M. Kameda, M. Hamada, and T. Takeuchi. 1997. Resormycin, a novel herbicidal and antifungal antibiotic produced by a strain of *Streptomyces platensis*. I. Taxonomy, production, isolation and biological properties. *J. Antibiot.* 50:1020–1025.
- Imbert, T. F. 1998. Discovery of podophyllotoxins. *Biochimie* 80:207–222.
- Jelenska, J., M. J. Crawford, O. S. Harb, E. Zuther, R. Haselkorn, D. S. Roos, and P. Gornicki. 2001. Subcellular localization of acetyl-CoA carboxylase in the apicomplexan parasite *Toxoplasma gondii*. *Proc. Natl. Acad. Sci.* 98:2723–2728.
- Jomaa, H., J. Wiesner, S. Sanderbrand, et al. 1999. Inhibitors of the non-mevalonate pathway of isoprenoid biosynthesis as antimalarial drugs. *Science* 285:1573–1576.
- Kastanias, M. A. and M. Chrysai-Tokousbalides. 2000. Herbicidal potential of pyrenophorol isolated from a *Dreschlera avenae* pathotype. *Pest Manag. Sci.* 56:227–232.
- Kati, V. and R. J. Froud-Williams. 1999. Studies on allelopathic potential of various cereal cultivars on selected test species. *Brighton Crop Prot. Conf., Weeds* 2:579–580.
- Kossanel, J. P., J. Martin, P. Anelle, M. Peinot, J. K. Vallet, and K. Kurnej. 1977. Inhibition of growth of young radicles of maize by exudates in culture solutions and extracts of ground roots of *Chenopodium album* L. Pages 77–86 in A. M. Grodzinsky, ed. *Interactions of Plants and Microorganisms in Phytocenoses*. Kiev: Naukova Dumka. [In Russian]
- Lang-Unnasch, N., M. E. Reith, J. Munholland, and J. R. Barta. 1998. Plastids are widespread and ancient in parasites of the phylum Apicomplexa. *Int. J. Parasitol.* 28:1743–1754.
- Lasceve, G. and F. Gaugain. 1990. Effects of usnic acid on sunflower and maize plantlets. *J. Plant Physiol.* 136:723–727.
- Lehnen, L. P., Jr., M. A. Vaughan, and K. C. Vaughn. 1990. Terbutol affects spindle microtubule organizing centres. *J. Exp. Bot.* 41:537–546.
- Lehnen, L. P., Jr. and K. C. Vaughn. 1992. The herbicide sindone B disrupts spindle microtubule organizing centers. *Pestic. Biochem. Physiol.* 44:50–59.
- Le Tourneau, D., G. D. Failes, and H. G. Heggeness. 1956. The effect of aqueous extracts of plant tissue on germination of seeds and growth of seedlings. *Weeds* 4:363–368.
- Liebermann, B., R. Ellinger, and E. Pinet. 1996. Isotontoxin, a conversion product of the phytotoxin tentoxin. *Phytochemistry* 42:1537–1540.
- Lin, W. X., K. U. Kim, and D. H. Shin. 2000. Rice allelopathic potential and its modes of action on barnyardgrass (*Echinochloa crus-galli*). *Allelopathy J.* 7:215–224.
- Lin, L.-J., G. Peiser, B.-P. Ying, K. Mathias, F. Karasina, Z. Wang, J. Itatani, L. Green, and Y.-S. Hwang. 1995. Identification of plant growth inhibitory principles in *Alisanthus altissima* and *Castela tortuosa*. *J. Agric. Food Chem.* 43:1708–1711.
- Lindon, J. C., J. K. Nicholson, and D. Ian. 1996. The development and application of coupled HPLC–NMR spectroscopy. *Adv. Chromatogr.* 124:315–382.
- Liu, D. L. and N. E. Christians. 1994a. Herbicidal activity of a hydrolysed corn gluten meal on three grass species under controlled environments. *J. Plant Growth Regul.* 13:221–226.
- Liu, D. L. and N. E. Christians. 1994b. Isolation and identification of

- root-inhibiting compounds from corn gluten hydrolysate. *J. Plant Growth Regul.* 13:227–230.
- Liu, D. L. and N. E. Christians. 1996. Bioactivity of a pentapeptide isolated from corn gluten hydrolysate on *Lolium perenne* L. *J. Plant Growth Regul.* 15:13–17.
- Liu, D. L. and N. E. Christians. 1997. Inhibitory activity of corn gluten hydrolysate on monocotyledonous species. *HortSci.* 32:243–245.
- Lucena, J. M. and J. Doll. 1976. Efectos inhibidores de crecimiento del coquito (*Cyperus rotundus* L.) sobre sorgo y soya. *Rev. Comalfi* 3:241–256.
- Lydon, J. and S. O. Duke. 1999. Inhibitors of glutamine biosynthesis. Pages 445–463 in B. K. Singh, ed. *Plant Amino Acids: Biochemistry and Biotechnology*. New York: Marcel Dekker.
- Macias, F. A., T. Ascension, J. M. G. Molinillo, R. M. Valera, and D. Castellano. 1996. Potential allelopathic sesquiterpene lactones from sunflowers leaves. *Phytochemistry* 43:1205–1215.
- Macias, F. A., J. M. G. Molinillo, J. C. Galindo, R. M. Varela, A. M. Simonet, and D. Castellano. 2001. The use of allelopathic studies in the search for natural herbicides. *J. Crop Prod.* 4:237–255.
- Macias, F. A., R. M. Varela, A. Torres, and J. G. M. Molinillo. 1999. Potential of cultivar sunflowers (*Helianthus annuus* L.) as a source of natural herbicidal templates. Pages 531–550 in Inderjit, K.M.M. Dakshiny, and C. L. Foy, eds. *Principles and Practice in Plant Ecology*. Boca Raton, FL: CRC Press.
- Mitchell, G., D. W. Bartlett, T. E. M. Fraser, T. R. Hawkes, D. C. Holt, J. K. Towson, and R. A. Wichert. 2001. Mesotrione: a new selective herbicide for use in maize. *Pest. Manag. Sci.* 57:120–128.
- Morré, D. J., P. A. Grieco, and D. M. Morré. 1998. Mode of action of the anticancer quassinoids—inhibition of the plasma membrane NADH oxidase. *Life Sci.* 63:595–604.
- Narwal, S. S. 2000. Weed management in rice–wheat rotation by allelopathy. *Crit. Rev. Plant Sci.* 19:249–260.
- Narwal, S. S., T. Singh, J. S. Hooda, and M. K. Kathuria. 1999. Allelopathic effect of sunflower on succeeding summer crops. 1. Field studies and bioassays. *Allelopathy J.* 6:35–48.
- Neale, M. 2000. The regulation of natural products as crop-protection agents. *Pest Manag. Sci.* 56:677–680.
- Netzly, D. H. and L. G. Butler. 1986. Root of sorghum exude hydrophobic droplets containing biologically active components. *Crop Sci.* 26:775–778.
- Netzly, D. H., J. L. Riopel, G. Ejeta, and L. G. Butler. 1988. Germination stimulants of witchweed (*Striga asiatica*) from hydrophobic root exudate of sorghum (*Sorghum bicolor*). *Weed Sci.* 36:441–446.
- Nimbal, C. I., J. Pedersen, C. N. Yerkes, L. A. Weston, and S. C. Weller. 1996. Activity and distribution of sorgoleone in grain sorghum germplasm. *J. Agric. Food Chem.* 44:1343–1347.
- Oliva, A., R. M. Moraes, S. B. Watson, S. O. Duke, and F. E. Dayan. 2002. Aryltetralin lignans inhibit plant growth by affecting the formation of mitotic microtubular organizing centers. *Pestic. Biochem. Physiol.* 72:45–54.
- Olofsdotter, M., D. Navarez, M. Rebulanan, and J. C. Streibig. 1999. Weed suppressing rice cultivars—does allelopathy play a role? *Weed Res.* 39:441–454.
- Pachlatko, J. P. 1998. Natural products in crop protection. *Chimia* 52:29–47.
- Paerl, H. W. and C. S. Tucker. 1995. Ecology of blue-green algae in aquaculture ponds. *J. World Aquacult. Soc.* 26:109–131.
- Parenti, L. R. and E. L. Rice. 1969. Inhibitional effects of *Digitaria sanguinalis* and possible role in old-field succession. *Bull. Torrey Bot. Club* 96:70–78.
- Patrick, Z. A. and L. W. Koch. 1958. Inhibition of respiration, germination, and growth by substances arising during decomposition of certain plant residues in soil. *Can. J. Bot.* 36:621–647.
- Patrick, Z. A. and L. W. Koch. 1963. The adverse influence of phytotoxic substances from decomposing plant residues on resistance of tobacco to black root rot. *Can. J. Bot.* 41:747–758.
- Pereda-Miranda, R., R. Mata, A. L. Anaya, D. B. M. Wickramaratne, J. M. Pezzuto, and A. D. Kinghorn. 1993. Tricolorin A, major phyto-growth inhibitor from *Ipomoea tricolor*. *J. Nat. Prod.* 56:571–582.
- Pezzuto, J. M., C. W. W. Beecher, H. H. S. Fong, et al. 1998. Recent studies on biologically active natural products. Pages 94–107 in Attar-Rahman and M. I. Choudhary, eds. *New Trends in Natural Products Chemistry*. The Netherlands: Hardwood Academic Publishers.
- Pillinger, J. M., J. A. Cooper, and I. Ridge. 1994. Role of phenolic compounds in the antialgal activity of barley straw. *J. Chem. Ecol.* 20:1557–1569.
- Pillmoor, J. B. 1998. Carbocyclic coformycin: a case study of the opportunities and pitfalls in the industrial search for new agrochemicals from nature. *Pestic. Sci.* 52:75–80.
- Pimm, S. L., G. J. Russell, J. L. Gittleman, and T. M. Brooks. 1995. The future of biodiversity. *Science* 269:347–350.
- Qasem, J. R. 1995. Allelopathic effects of *Amaranthus retroflexus* L. and *Chenopodium murale* L. on vegetable crops. *Allelopathy J.* 2:49–56.
- Renner, S. S. and R. E. Ricklefs. 1998. Herbicidal activity of domatia-inhabiting ants in patches of *Tococa guianensis* and *Clidemia heterophylla*. *Biotropica* 30:324–327.
- Rice, E. L. 1964. Inhibition of nitrogen-fixing and nitrifying bacteria by seed plants. *I. Ecology* 45:824–837.
- Rice, E. L. 1984. *Allelopathy*. Orlando, FL: Academic Press. 422 p.
- Rice, M. J., M. Legg, and K. A. Powell. 1998. Natural products in agriculture—a view from the industry. *Pestic. Sci.* 52:184–188.
- Rimando, A. M., F. E. Dayan, M. A. Czarnota, L. A. Weston, and S. O. Duke. 1998. A new photosystem II electron transfer inhibitor from *Sorghum bicolor*. *J. Nat. Prod.* 61:927–930.
- Rimando, A. M., F. E. Dayan, J. R. Mikell, and R. M. Moraes. 1999. Phytotoxic lignans of *Leucophyllum frutescens*. *Nat. Toxins* 7:39–43.
- Rimando, A. M., M. Olofsdotter, S. O. Duke, and F. E. Dayan. 2001. Searching for rice allelochemicals: an example of bioassay-guided isolation. *Agron. J.* 93:16–20.
- Roberts, R., C. W. Roberts, J. J. Johnson, et al. 1999. Evidence for the shikimate pathway in apicomplexan parasites. *Nature* 393:801–805.
- Rojas, I. S., B. Lotina-Hennsen, and R. Mata. 2000. Effect of lichen metabolites on thylakoid electron transport and photophosphorylation in isolated spinach chloroplasts. *J. Nat. Prod.* 63:1396–1399.
- Romagni, J. G., S. O. Duke, and F. E. Dayan. 2000a. Inhibition of plant asparagine synthetase by monoterpene cineoles. *Plant Physiol.* 123:725–732.
- Romagni, J. G., G. Meazza, N. P. D. Nanyakkara, and F. E. Dayan. 2000b. The phytotoxic lichen metabolite, usnic acid, is a potent inhibitor of plant *p*-hydroxyphenylpyruvate dioxygenase. *FEBS Lett.* 480:301–305.
- Saxena, S. and A. K. Pandey. 2001. Microbial metabolites as eco-friendly agrochemicals. *Appl. Microbiol. Biotechnol.* 55:395–403.
- Schrader, K. K., F. E. Dayan, S. N. Allen, M. Q. de Regt, C. S. Tucker, and R. N. Paul, Jr. 2000a. 9,10-Anthraquinone reduces the photosynthetic efficiency of *Oscillatoria perornata* and modifies cellular inclusions. *Int. J. Plant Sci.* 161:265–270.
- Schrader, K. K., M. Q. de Regt, P. D. Tidwell, C. S. Tucker, and S. O. Duke. 1998a. Compounds with selective toxicity towards the off-flavor metabolite-producing cyanobacterium *Oscillatoria* cf. *chalybea*. *Aquaculture* 163:85–99.
- Schrader, K. K., M. Q. de Regt, P. R. Tidwell, C. S. Tucker, and S. O. Duke. 1998b. Selective growth inhibition of the musty-odor producing cyanobacterium *Oscillatoria* cf. *chalybea* by natural compounds. *Bull. Environ. Contam. Toxicol.* 60:651–658.
- Schrader, K. K., S. O. Duke, S. K. Kingsbury, C. S. Tucker, M. V. Duke, C. P. Dionigi, D. F. Millie, and P. V. Zimba. 2000b. Evaluation of ferulic acid for controlling the musty-odor cyanobacterium, *Oscillatoria perornata*, in aquaculture ponds. *J. Appl. Aquacult.* 10:1–16.
- Schrader, K. K. and M. D. Harries. 2001. Compounds with selective toxicity toward the musty-odor cyanobacterium *Oscillatoria perornata*. *Bull. Environ. Contam. Toxicol.* 66:801–807.
- Schumacher, W. J., D. C. Thill, and C. A. Lee. 1982. The allelopathic potential of wild oat (*Avena fatua* L.) on spring wheat (*Triticum aestivum* L.) growth. Urbana-Champaign, IL: North American Symposium on Allelopathy, November 14–17, 1982. [Abstract]
- Singh, S. P. 1968. Presence of a growth inhibitor in the tubers of nutgrass (*Cyperus rotundus* L.). *Proc. Indian Acad. Sci.* 67:18–23.
- Sugawara, F. 2000. Phytotoxins as potential herbicides. Pages 113–128 in S. S. Narwal, R. E. Hoagland, R. H. Dilday, and M. J. Reigosa, eds. *Allelopathy in Ecological Agriculture and Forestry*. Amsterdam: Kluwer Academic Publishers.
- Tanaka, T., T. Fukushima, Y. Tsujino, and T. Fujimori. 1997. Nigrosporins A and B, new phytotoxic and antibacterial metabolites produced by a fungus *Nigrospora oryzae*. *Biosci. Biotech. Biochem.* 61:1848–1852.
- Tanaka, T., K. Hatano, and M. Watanabe. 1996. Isolation, purification and identification of 2,5-anhydro-D-glucitol as a phytotoxin from *Fusarium solani*. *J. Nat. Toxins* 5:317–329.
- ten Kate, K. and S. A. Laird. 1999. *The Commercial Use of Biodiversity. Access to Genetic Resources and Benefit-Sharing*. London: Earthscan Publications. pp. 13–33.

- Tinnin, R. and C. Muller. 1971. The allelopathic potential of *Avena fatua*: influence on herb distribution. *Bull. Torrey Bot. Club* 98:243–250.
- Tucker, C. S. and S. W. Lloyd. 1987. Evaluation of potassium ricinoleate as a selective blue-green algicide in channel catfish ponds. *Aquaculture* 65:141–148.
- Unruh, J. B., N. E. Christians, and H. T. Homer. 1997a. Herbicidal effects of the dipeptide, alaninyl-alanine, on perennial ryegrass (*Lolium perenne* L.) seedlings. *Crop Sci.* 37:208–211.
- Unruh, J. B., N. E. Christians, and H. T. Homer. 1997b. Mitotic and ultrastructure changes in root meristems of grass seedlings treated with alaninyl-alanine. *Crop Sci.* 37:1870–1874.
- Van Aller, R. T. and G. F. Pessoney. 1982. USM algal research team makes major off-flavor/water quality discovery. *Aquacult. Mag.* 8:18–22.
- Van Aller, R. T., G. F. Pessoney, V. A. Rogers, E. G. Watkins, and H. G. Leggett. 1985. Oxygenated fatty acids: a class of allelochemicals from aquatic plants. *ACS Symp. Ser.* 268:387–400.
- van der Ploeg, M., C. S. Tucker, and C. E. Boyd. 1992. Geosmin and 2-methylisoborneol production by cyanobacteria in fish ponds in the southeastern United States. *Water Sci. Technol.* 25:283–290.
- Vaughan, M. A. and K. C. Vaughn. 1988. Mitotic disrupters from plants and their potential use as herbicides. *Weed Technol.* 2:533–539.
- Vaughan, M. A. and K. C. Vaughn. 1990. DCPC causes cell plate disruption in wheat roots. *Ann. Bot.* 65:379–388.
- Vaughn, S. F. and M. A. Berhow. 1998. 1-Cyano-2-hydroxy-3-butene, a phytotoxin from crambe (*Crambe abyssinica*) seedmeal. *J. Chem. Ecol.* 24:1117–1126.
- Vaughn, S. F., R. A. Boydston, and C. A. Mallory-Smith. 1996. Isolation and identification of (3-methoxyphenyl)acetonitrile as a phytotoxin from meadowfoam (*Limnanthe alba*) seedmeal. *J. Chem. Ecol.* 22:1939–1949.
- Velu, G. and R. Aruna. 1996. Allelopathic impact of purple nutsedge (*Cyperus rotundus*) and bermuda grass (*Cynodon dactylon*) on soybean (*Glycine max*). *Indian J. Agric. Sci.* 66:363–365.
- Waller, G. 1987. Allelochemicals: Role in Agriculture and Forestry. Washington: ACS Symposium Series 330. 606 p.
- Welch, I. M., P. R. F. Barrett, M. T. Gibson, and I. Ridge. 1990. Barley straw as an inhibitor of algal growth I: studies in the Chesterfield Canal. *J. Appl. Phycol.* 2:231–239.
- Weston, L. 1996. Utilization of allelopathy for weed management in agroecosystems. *Agron. J.* 88:860–866.
- Weston, L. A., C. I. Nimbai, and P. Jeandet. 1999. Allelopathic potential of grain sorghum (*Sorghum bicolor* (L.) Moench) and related species. Pages 467–478 in Inderjit, K.M.M. Dakshiny, and C. L. Foy, eds. *Principles and Practice in Plant Ecology*. Boca Raton, FL: CRC Press.
- Wibiwo, D. N. 1996. Effect of root and shoot extracts of purple nutsedge (*Cyperus rotundus* L.) at different concentrations on root nodule formation, growth, and yield of soybean (*Glycine max* (L.) Merr.). Pages 139–147 in *Proceedings of the Symposium on Biology and Management of Weeds and fourth Tropical Weed Science Conference, 1994*. Bogor, Indonesia: SEAMEO BIOTROP Special Publication No. 58.
- Williams, J. T. 1964. A study of the competitive ability of *Chenopodium album* L. *Weed Res.* 4:283–295.
- Willis, R. J. 2000. *Juglans* spp. juglone and allelopathy. *Allelopathy J.* 7:1–55.
- Wills, G. D., C. S. Tucker, and E. J. Jones. 1999. Effect of barley straw for the control of off-flavor in pond-raised catfish. *Proc. South. Weed Sci. Soc.* 52:227–230.
- Yoneyama, K., T. Saruta, M. Ogasawara, M. Konnai, T. Asami, T. Abe, and S. Yoshida. 1996. Effects of grandinol and related phloroglucinol derivatives on transpiration and stomatal closure. *Plant Growth Regul.* 19:7–11.
- Young, C. C. 1986. Autointoxication of *Asparagus officinalis* L. Pages 171–188 in A. Putnam and C. S. Tang, eds. *The Science of Allelopathy*. New York: Wiley Interscience.
- Zeidler, J., J. Schwender, C. Muller, J. Wiesner, C. Weidemeyer, E. Beck, H. Jomaa, and H. K. Lichtenthaler. 1998. Inhibition of the non-mevalonate 1-deoxy-D-xylulose 5-phosphate pathway of plant isoprenoid biosynthesis by fosmidomycin. *Z. Naturforsch.* 53c:980–986.

Received November 13, 2001, and approved November 13, 2001.